

# BIOGENESIS III™ SD

## SEED COATING MATERIAL

### APPLICATIONS AND DIRECTIONS:

#### PLANTER BOX:

Biogenesis IIISD can be applied dry at a rate of 2 ounces per 100 pounds of seed in drill box.

#### SLURRY:

Can be applied at a rate of 1 oz per 100 pounds of seed, mixed in liquid form. The material should be used within 4 hours of mixing. When product has dried on the seed, seed can be stored as normal with very little microbe deterioration.

Net Weight: \_\_\_\_ gm.

Exp. Date: \_\_\_\_\_

Lot #: \_\_\_\_\_



**TAINIO TECHNOLOGY  
& TECHNIQUE, INC.**

*Natural Growing Through Technology*

S. 12102 ANDRUS RD., CHENEY, WA 99004

[www.tainio.com](http://www.tainio.com)

### CONTENTS:

Live soil microbes, including:  
5 strains of *Arthrobacter*  
*Azospirillum lipoferum*  
2 stains of *Azotobacter*  
5 strains of *Bacillus*  
2 strains of *Bacteroides*  
*Kurtha zopfil*  
3 strains of *Pseudomonas*

### CAUTION:

Keep out of reach of children and pets.  
Do not swallow.

### CONTAINS LIVE SOIL MICROBES:

Store below 120 degrees F.  
Keep out of direct sunlight.